



VACCINE	DOSES	AGE FOR ADMINISTRATION
BCG Vaccine	1 DOSE	AT BIRTH (OR AS EARLY AS POSSIBLE UP TO 1 YEAR)
Hepatitis B	4 DOSES	AT BIRTH, 6 WEEKS, 10 WEEKS, AND 14 WEEKS
Oral Polio Vaccine	4 ORAL + 1 IPV	AT BIRTH, 6 WEEKS, 10 WEEKS, 14 WEEKS, AND 16-24 MONTHS (BOOSTER)
DPT vaccine	5 DOSES	6 WEEKS, 10 WEEKS, 14 WEEKS, 16-24 MONTHS
Hib Vaccine	3 DOSES	6 WEEKS, 10 WEEKS, AND 14 WEEKS
Rotavirus Vaccine	3 DOSES	6 WEEKS, 10 WEEKS, AND 14 WEEKS
Pneumococcal Vaccine	3 DOSES + 1 BOOSTER	6 WEEKS, 10 WEEKS, 14 WEEKS, AND A BOOSTER AT 12-15 MONTHS
MR vaccine	2 DOSES	9-12 MONTHS AND 16-24 MONTHS
JE Vaccine	2 DOSES	9-12 MONTHS AND 16-24 MONTHS
Vitamin A	9 DOSES	FIRST DOSE AT 9 MONTHS, THEN EVERY 6 MONTHS UP TO 5 YEARS



VACCINE

DOSES

AGE FOR ADMINISTRATION

HPV Vaccine

3 DOSES

9-14 YEARS (TWO DOSES, 6 MONTHS APART) OR ≥15 YEARS
(THREE DOSES: 0, 1, 6 MONTHS)

Td vaccine

BOOSTER EVERY 10 YEARS

10 YEARS, 16 YEARS, AND EVERY 10 YEARS THEREAFTER

Influenza Vaccine

ANNUAL DOSE

ANNUALLY FOR ALL AGES, ESPECIALLY HIGH-RISK GROUPS

Pneumococcal Vaccine

1 OR 2 DOSES + BOOSTER

ADULTS ≥65 YEARS, OR EARLIER FOR INDIVIDUALS WITH
CHRONIC CONDITIONS

Hepatitis A

2 DOSES

AT LEAST 6 MONTHS APART, ESPECIALLY FOR THOSE AT HIGH
RISK OF INFECTION

Hepatitis B

3 DOSES

0, 1, AND 6 MONTHS FOR ADULTS NOT PREVIOUSLY VACCINATED

Shingles Vaccine

2 DOSES

FOR ADULTS AGED 50 YEARS AND OLDER

